

# Low dose diuretics for treatment of advanced schistosomiasis cirrhotic ascites after regression Prevention of recurrence clinical observations

ZhuJijun<sup>1</sup>WeiDongping<sup>1</sup>, ShenHongxia<sup>2</sup>

<sup>1</sup> 432100LakeNorthfilialsenseCitybookyangsoilAreaHealth Serviceheart

<sup>2</sup> 432100Xiaogan Xiaonan District Wolong TownshipBlood Station2

**Abstract:** objective:theclinical Observa LionofRecmtenceprevention oi'smallDsystemeofdiuretictratarlvance,lschist osemianameminCir

RHo♀Isascitessushi.-le.-1.metho.-ls:collectingneWAdvancedschileafSunomiasisCIRRhbrushesisasone  
cites patientS.fromTo2013,to the PA  
tientsTheAskmptomatictreatmentat thesaMeTime,continuouLeafFlyForalongtimelogIve  
peopleMallDotoeofDilu-etic caneffectIvelyPreventarecmtenceof arlvanced schistPlayMiasiscit  
RhosI♀Ascitesmjnthelaterplaysaveryimportantrole inClinical application.

**Keywords:** SmAll. -landofProducturetic:a.-1-

vancedliver,hIaddress.miasis:CinhoPeopleis withawithites:recunencePrevention

Schistosoma japonicum is a threat to our peasant bodythe most serious parasitic disease in health.is sucked by Japanese blood.worm intointo body.and parasitic on the portal vein system.causesonespecies Chishengfrom.Human infection in Japanschistosomiasis is mainly in contact with the skin of Schistosoma japonicumthetail of the water has a directRelationship.Infected with Japanese bloodThe main lesions of the disease were imprinted by the worm in the liver andinthe colon, and the granuloma.,IProThe beds are acuteperiod.slowand late.The main manifestation of acute phase isfever.Allergy Skin should,abdominal symptom; chronic period withLiver and spleen enlargement mainly; Late major lesions are portal veinPeripheral fibrosis.The final can be left for portal hypertensionsyndrome,splenomegalyandliver cirrhosis ascites.Advanced schistosomiasis

(withlateblood) patients with cirrhosis of ascites undergo a pairofFade After treatment.However, it is often easy tosend.CausesThe severity of the patient to worsen,,My center uses long interruptedTreatment of late-stage blood suction with small doses of furosemide and endoscopic therapydisease of cirrhotic ascites after regressionpatient.Found liversclerosing ascites recurrence rateMingnexttwo

Information and methods

2010-2013Annual New Night Blood (ascitestype) treatmentPatient basic condition.Seetablel.

late blood cirrhosis ascites.Diagnostics and dischargestandardreference thetechnical scheme for the treatment of patients withadvanced schistosomiasis in Hubei Provincesection,

Method:to patients with a diuretic,patch no eggswHITE,SecurityLiveretextsyndromecuretherapy ascites.Ascites subsidedafterfollowingRenew

servicewithfurosemidelomg.11knowDayandsnailinsidevinegar20mg.1Times/Day.15daysstopwithlWeek.thenfollowingc  
ontinue taking.,outhouseafterlongperiodKennedyholding servicewith.Nextevery3Month set  
periodcheck.CheckCheckLiver.kidneyworkcan,abdomensectionBSuper, and so on=

Copyright © 2017 Author(s).

This is an open-access article distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 International License (<http://creativecommons.org/licenses/by-nc/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Knot Fruit

yearpartLyearinsideBellyWaterRecurrence.Seetable2.,,

years/year inside ascites recurrence Analysis. See table 3.,,

To ask the

Ascites is formed by a combination of factors which causes the water to remain in the abdominal cavity of the main reason is portal high pressure. The internal dirty blood vessel dilation causes the effective blood volume matrix to be the main factor that causes water retention and ascites formation and aggravation [11,], from 4 years treatment of New Left patients observation results, each year {sub-abdomen water recovery left rate %} around and 6 months after occurrence. The application of small dosage diuretic Liu to control ascites recurrence is obvious. Large amount foreign research report road. In the juice: count have ascites or ascites-free liver hard to free water in abdominal cavity. The patient's body was found to be delivered to the kidney, the rate of the distal end of the tubules clear show decrease. This indicates that the system of equal electrolytes in the renal small tube near the heavy absorbance increase, and kidney small distal tube W is significant

The decrease is caused by free water in the patient's body.

cause heart Even if the patient does not have a combined ascites. Kidney small

small do kidney f diuretic therapy. Results: each

Fly Ear a fly ear ascites REC Unmilesnce is as follows.

Table 1 2010-2013 Year new rub Night blood (ascites type) basic conditions for treatment of patients

Year Cases Man Woman year Ling average year Ling

In 2010, a total of 5 cases of Frencho-rence (31.25%), after R June 1 ca Mourning after Sep

ember 2 Cases. After December 2 cases: LN%, a total of 4 cases of Frencho-

rence (26.67 %), after June 1 ca clear after SEP

Temher 2 cases. After December 1 Ca Goodes: LN 2012. A total of 3 Yang back Ran Torecun: ence

(30%). After June 1 Ca back to "after September er" 2 cases people: in 2013 a total of 3 cases of Frencho-rence What

(30%), after June 1 ca where, After September 1 Ca only yes. After December 1 cases. Conclusion:

Long-term, continuous with application

Table 2 Prepare year 1 Recurrence of ascites during the year (example) table 3 Prepare year 1 ascites Recurrence Analysis during the year (%)

Year Total 3 Months. Months 9 Months mouth months Afternoon Total 3 Month. Month 9 month Month

Application of two-hole laparoscopic cholecystectomy in primary hospital

Guou

650000 hepatobiliary Surgery, Yunnan Milei County People's hospital

Doi: Ten.3969/j.issn.1007 614x.2013.24.

Pick to Mesh: share two.: JI, method anti-endoscopic gallbladder resection (LC) in clinical work should with experience, and analyzing it should with price value method: on the 15 SE example two.: JI, law LC Profiling summary, results: two.: JI, method LC and General 3 hole

comparison in operation break, Surgery medium out Blood, hospitalization time, hospitalization fee with, ill feeling recovery, etc. difference meta-statistical significance. knot theory: in the grass-roots hospital can be carried out according to the patient's request two.: JI, method LC, but not recommended for general surgery.,

keywords: JI, laparoscopy month Dan Atlas resection grass roots Medical Courtyard Apply Value

Abstract:

objective: we want to also share the experience of Two hole LC in clinical work. - 1 analyze its application value. Methoonels: 158 ca back softwo hole LC In We Hospi Talw As

a NALyzed and summa Port Z Brush Onel. results: The two

hole LC method and The conventional Three hole

LC method Is compared in operation Time bleeding

d Ming operation Hospitaliza Lion Time, Cos Tand ill ness recover v and there was no significant Dit forence. CONCLUS. ion: we can According to the requirements of PA

Show LC. You have mastered the: Operational Skills Operation. With four-hole method, three-hole method and two-hole method succeeded for Patient 2000 cases. Case and using the regular to gauge laparoscopic devices try to execute multiple cases via umbilical Chanyu Laparoscopic cholecystectomy Port present summary year Current two-hole method LC case 158 Example. Analysis of its clinical application at the primary level of the Medical hospital plant. The report is as follows

#### Information and methods

This group of patients 1% Example. man example. Women the example. Year age old. average old. where slow sex gall the sac inflammatory gallbladder stone 105 example. gall saciflesh Change Example. urgent sex gall capsule inflammatory close and Stone one example with accent LJ Palace Production History 5 example. appendectomy History 3 example. combined with hypertension example, Merge Diabetes 8 example I have patients routinely check abdomen after admission B Super, ECG Figure, X chest x, blood routine and blood biochemistry, Blood coagulation can wait. when patients with acute cholecystitis and calculus and 2 example sleep cholelithiasis patients with biliary duct mild enlargement

Zhang Fly slightly elevated bilirubin. fr MRCPT Check excluding bile duct and membranous gland lesions. Acute gallbladder Inflammatory and calculous patients are treated with a silent inflammation. the symptom is well verified go after 23 Day Hand operation L""

Surgical Side method: all take gas pipe intubation body linen Drunk. Do not place the urinetube before surgery. There is a stomach inside expansion gas Obviously affects the retention of the stomach tube. Take navel upper margin do 10 11 11 11 arc shape cut Port. wear string set into Lomm

Explorer Location Relationship. with electric hook horizontal form knife opening neck and calot Blue corner serous membrane. Use the separator to lower the size of the film and its adipose tissue. Clearly reveal the gallbladder tube, ij\_\$. Explorer. Liver total tube and gallbladder artery. relaxed traction shop. toi\_\$. Total Top of pipe 0.5 cm use Bio-folder folder Ask the cystic duct. In the top of it with an iron clamp | -hole gallbladder tube. cut between two clips break gall duct. Use the raw object clip to close the gallbladder artery and cut break Mountain Open gallbladder serous membrane and Record Tl from gallbladder liver bed to complete under knife gallbladder = electrocoagulation gallbladder bed stop bleeding. depending on whether or not the indwelling drainage is determined by the operation,,, enlarged navel jab main remove gallbladder. such as Cholecystitis water swollen heavier or larger stones. you can also cut and close the two incision to make it merge to one cut. The gall sac is removed.

#### Knot Fruit

158 example patients with 1 cases due to obesity, 2 Example because of a serious difficulty in satisfying exposure and operation. Change The three hole method to complete in 3 The remaining cases were performed with two-hole methods successfully completed,, all cases no yellow plague after operation. high hot. Abnormal abdominal pain shape! "1 Patients with fewer blood infiltration. 2 example inflammation more heavy, patient Yu Jian sudden indwelling Abdominal search | flow tube. after 12 Day Unplugged. Primer Flow tube No further bleeds and bile outflow,,, Surgery Time

minutes. Average on minutes, intraoperative bleeding

31}. average 1 Mountain sword Turkistan under wound pain Pain Example. Navel Create mouth pain Pain 8 example. All mild

tient people In Primary Hospital people To carry out the 2

Trocar. establish CO<sub>2</sub> pneumoperitoneum. Pressure to maintain ten ~ pain tolerable. Special handling. is not 3 6 days

Hole method LC, buttoh onk! Not he extend brushes One to then onnal Operation mo'onele

Keyword two hole Laparoscopic choleHtontectomy: basic Lev El when pit Al: application to value

Laparoscopic Gall SAC Excision (LC) now widely shipped with on clinical. with its wound small, restore quick, and so on Benefits received by Doctor and patient by Li home from 1998 Year Open

12 11 11 11 Hg. Set into 30° laparoscopy. observe the internal organs of the abdominal cavity and the conditions of the

retina and the abdominal wall 3 5 cm to bypass the sickle ligament under the lens in its right side. Wear Stab number of people 10111111 Trocar as the master as a hole. On the left side of an arc cut on the navel about 0.5 cm. The places at tattoo 5111111 trocar. Set people Bullets Spring traction pliers or detach Clamp,, tilt to the left of obese people 30. Corner and Fetch header high droptwo lifting gall with traction pliers sac body tone fl show Dew Gall SAC Neck. Clear Gall sac tube, Gall Mains and Liver

Better; off-bed activity time 4 [hours. average 7 small when; all after operation 6 Hour to to drinking. to eat a low-fat, semi liquid diet the next day after surgery; in hospital 37 Day. Yu Jun 5 day,, After surgery 3 month. six months and 1 Year follow-up. Success Visit 156 example. 2 after Operation 2 fabric with upper abdomen mild pain. check B not see exception give oral drug treatment better. 1 example after 1 year re-send Biletube stone. 1 Example after year duplicate check B Super. Liver work or MRCP all not see bile duct narrowly

The obstruction of the distal end of the tube directly affects the drinking water with this the body's diuretic. Benefits System effect. Guide effect decrease I'll. so. Long-term intermittent use of small doses of diuretics can reduce the amount of uranum retention in patients with liver cirrhosis. Prevention of Liver sclerosis ascites complex send. patients with liver cirrhosis can produce health large amount rates quality stress Harmony Solid States same as. These two excitement element can induce the occurrence of ascites Port Water na? Green stay is liver

Harden the original friend change. Ascites is a node of liver cirrhosis fruit J Long-term intermittent use of small dose diuretics improves Water Retention. Fundamentally improve the liver hard. Diuretic agent can play a role in the internal glare of the state of the body.. can also enhance the drainage effect of the kidneys. To maintain the hydro-electrolyte balance., intermittent take ease of suffering on diuretic tolerance by to up to most good effect fruit 3

## References

---

1 New!!! U Forest Fruit is. Real with inside Science [M] Beijing: People's Guardian Health publication Note. 2009  
 2 wait J Violin Hallyuan., and so on. regression of cirrhotic ascites withi with overall inside electrolyte Balance | Self. study research [JJ. Chinese Digest Miscellaneous Log. 19922 (2): 84-86.