

sendfertilityNavigationoff section dislocationtreatmentmedium bonefoldpreanti

Lou Smart EnglishALily

Guiyang MCH Hospital pediatric orthopedics, Guizhouguiyang0 003

Middle Chart category number:R473.6document identifier:Barticle number:1000744X (2017)three block01

Abstract: Developmental Russian JointJHyatt(DOH) is the most common limb attachment for children, governingtreatmentvaries by age and the degree of pathological changes in the Russian joint.We have13 year8Month to2015year6Month opening operation in our hospitaldoHPost-operative raw femurfracturei1tline analysis and2015year6months to16Year6Month forDOHThe reason forThe recovery of the post after surgery, take humanity toBootstrap primaryComprehensive training and preventive measures to get good results.Thereport is as follows.

Keywords:Friend-fertile bony dislocation;Fracture;Prevent

1. Data and Methods

1.1 Onedata collect our homes2013year8months to2015year6MonthAdmission the development of the dislocation of the Russian joint is complete236Examplehungry), after treatment fracture of femur in process8Example(2.6%).age from2.5old to8Year,Ping all3.5Age, male3* * A, a., female5example.2example occurs on joint strong;after Operation8 months,5The example occurs after the Operation3 5a monthwhere1Example Direct force salt damage,5 example Fall Indirect injury,2example damages in functionaltraining.2015year6months to 2016year6month157Example(203do),take humanistic guidance for the main comprehensivetraining measures,scraping parents ' awareness of the disease and how to cooperatewith it,HVHousePre-trained squaremethod,discharged to perform joint functional training for the main "comprehensive"guide.

1.2 Precautionary method(1)rating Parentsdegree of education and family values,multiple previous toloyaltyChildand parents to introduce the purpose of the Operation,effect,postoperativecomplicationsand important for postoperative functional training,trainer jumps scientific Russian joint functiontraining.withchildrenperform lower limb muscles on the bed from the start of tractionJJand Bone hardness trainingpractice,and the most effective approach is the long muscular contraction training.themethod of training a feature is usedin thepicture'sformtoparents' show,practice in kind.(2)about affected version of activityComment,after the operation to observe loyalty son have Yuan hungry knuckle flexionMountainLTwinshrink,If standingLumbar Spine Pastextension,Pelvic front{?,Walk with shorter stride lengthrigidity step[I,based on rating results personalityturnssystemsettrainingtrainerDraw,trainingPracticedtheprocessr!1 Watch out forChildinverseshould,Action LightsoftSlowslow,with rhythmsex,in a certainBendStretchin-range step-by-Step,force size to sufferchildtolerable degree,prohibituse violence.(3) extranet setperiodbetweenthemuscular contraction trainingpracticefor the movement of the outer joint and the limb of the patient is not defined.Remove stonedo not have negative cocoon training after,throughXLline showshowboneretroactivelylivelonggood OK,Start weight Training[2)daily Turn offsection feature trainingpractice,3times,everytimetrainingBefore practicing infra-

Copyright © 2019 Author(s).

This is an open-access article distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 International License (<http://creativecommons.org/licenses/by-nc/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited

redZtype treatmentlight irradiation or local hot compressmin,I start trainingourwithCPMSecondary

to main, by a fact to become suffering child active motion to main, nurses and parents at the side assist with protect with child, avoid injury. (4) patients with poor active compliance, only. Line Passive training, enhancing communication with children, patiently cited guide, with suffer child interested. Tours Play, Mobilizing children main move parameter with, Product with rehabilitation training Extreme, during Game Towel Complete Training. you can also let children watch TV while training distracting force, Remove tension thread, to avoid excessive joint muscle tension. (5) vitamins * Oral "million U, monthly 1 times, used with 2 times. Calcium gluconate Port

Service Liquid 10 mL, every day 1 times. Add a lifeline and high calcium intake, Bask in the sun as much as possible. (6) Establish back visit system degree, via phone, Micro letter or network, etc. with the parent-connected Department, urging parents to suffer child Perform the function training practice, and urge parents to post-operation 1, 3, 6 month and 1 year with patient child come home complex check, to Master with Child recovery and problems in a timely manner.

A Communication as, -ai L, 348106129 QQ, C. M

2. knot Fruit

to 15 year 6 months to 20 Year 6 Month discharged 157 Example (203 Mirror) suffers from the child escapes with visit, femur occurs 'Self-folding L Example (0. Total %), age 3.9 old, women Sex, inside Network Fix LL postoperative 1 week Fall injury. the remainder is special.

3. Ask comments

the incidence of developmental dislocation of the lam joint is 3.8% 0, net report new live child The incidence is approximately 4% 0 10% 0 [3]. The important measure to prevent joint stiffness is work can be trained, in the process of functional training is very easy to cause the anger joint instability and the occurrence of forging

Remove the joints bit, and another complication, unfeature recovery period and training process

loyal side femoral fracture. Retrospective Analysis of these cases in our hospital one injury bed and imaging funding

Material Is; original because of can To: (1) postoperative yue Joint stiffness reason with hand Create Injury, after Operations solid when between LYao, causes articular Mandarin, muscle meat twin Shrink and joint ii \

Straight. "• J", All of our older loyalties are from rural remote areas Area Home courtroom, Parent

lack of knowledge of the disease knowledge, Bucket then ask your family to help them with functional exercise. recognized difference, make! \ Easy concurrent pin ankylosis, J and the child walk spay appears gait not stable, prone to falls cause fracture. (2) femur 'merchant' Density change: This group 8 Example DOH children with femoral fracture after recovery period, Femur X line level Slice tip osteoporosis. has the report way "• Iskeleton and inside "Pavilion Use of fixed steel rice can be because loyalty ni IJ bone density drop of limb low, loss of bone because bone terror intensity significantly under drop, Welcome to a smaller force Impact Mesh, can cause bone fold, This main H I all have femur internal fixation. (3) lack of scientific forging practice: In this group there are JJ; TJ example occurs during the enter line function Exercise 'mortar fold, analyzing its reason is failed to communicate well before loyal children. Loyal Children in the course of the exercise because of pain and fear, does not work with exercise and severely resists production live reaction, makes it harder to exercise

Large, unexpected bone occurrence fold back. Section loyalty, child joint function inferior to other similar children restore good, Parents and caregivers are in a hurry during the exercise in the Hope's heart, No analysis is made based on the specific conditions of the child during exercise, early shipping move before move large. (4) Improper home care causes accidental injury harm: regardless of hand surgery with non-surgical treatment DO Hall to network and restrict lower limb activities I paragraph between, when solution | Shi parents rush to get children to carry weight training, but child flat Balance Ability Low, walk not stable, It is very easy to fall or be broken by a cocoon. The femoral fracture. to increase parental attention to the degree, help children and parents recognize the disease, cure procedure Towel get stuck difficult, mobilizing children to participate actively, cooperate with rehabilitation training practice at the same time, also avoid accidental injury. Yang

References

1. Zhengcolore,LeeshowCloud.RealwithnetprotectionLearn[M].page1edition.PeopleGuardianliveoutversion FungL.:20101-110.
2. Sunlide,ShaoyiDevelopmental Russian Joint removalbitdiagnosisbreak[] cure [M.JinanSouth Publishing Fung1.:2002,170-176.
3. Drain the truemonth1Pediatric OrthopaedicsLearn(M).2EditionPeople's Health Mountain editionAgency,2005,77-93.
4. lagodd,kingFreeze Ice,ZhangMingang,etcDevelopmental pindislocationafterjGlare'lie'fromsecretLBranch 'sChangeandProBedmeaningsemantic[J].ChinaSmallPediatics Miscellaneouslog,201L38-6 1.
5. Li set, LeeJayLeeMing,etcDOHafterearlyPeriod RehabilitationTrainingPracticeFruitViewWatch[J].Mountain EastMedicalMedicine,2010,50 (22),86.
6. Xustate He.preoperativeRehabilitation Teaching EducationinallRussiaoffsectionpermutations[J].MediumcountrybonevsJointMiscellaneousLog,2014,3(9),3-655.